



## Risk assessment of *Salmonella* in Danish meatballs produced in the catering sector

Møller, Cleide Oliveira de Almeida; Nauta, Maarten; Schaffner, Donald W.; Dalgaard, Paw; Christensen, Bjarke Bak; Hansen, Tina Beck

*Publication date:*  
2015

[Link back to DTU Orbit](#)

### *Citation (APA):*

Møller, C. O. D. A. (Author), Nauta, M. (Author), Schaffner, D. W. (Author), Dalgaard, P. (Author), Christensen, B. B. (Author), & Hansen, T. B. (Author). (2015). Risk assessment of *Salmonella* in Danish meatballs produced in the catering sector. Sound/Visual production (digital)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

## International Journal of Food Microbiology

journal homepage: [www.elsevier.com/locate/ijfoodmicro](http://www.elsevier.com/locate/ijfoodmicro)



### Risk assessment of *Salmonella* in Danish meatballs produced in the catering sector



Cleide O. de A. Møller<sup>a</sup>, Maarten J. Nauta<sup>a</sup>, Donald W. Schaffner<sup>b</sup>, Paw Dalgaard<sup>c</sup>,  
Bjarke B. Christensen<sup>a,1</sup>, Tina B. Hansen<sup>a,\*</sup>

<sup>a</sup> National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark

<sup>b</sup> Department of Food Science, School of Environmental and Biological Science, Rutgers University, 65 Dudley Road, Food Science Building, Room 207, New Brunswick, NJ 08901-8520, USA

<sup>c</sup> National Food Institute, Technical University of Denmark, Søtofts Plads, Bygning 221, DK-2800 Kgs. Lyngby, Denmark


### Correspondence

Cleide Oliveira de Almeida Møller

E-mail: [clemo@food.dtu.dk](mailto:clemo@food.dtu.dk)

DTU Food

National Food Institute




# Outline:

- Introduction
- Objectives of the study
- Summarizing the performed work
- Process to build up the developed model
- Results
- Remarks and future perspectives

# Introduction

- *Salmonella* is a critical pathogen (CDC, 2011; EFSA, 2010).
- Pork still is an important source of salmonellosis (EFSA, 2010; van Hoek *et al.*, 2012; Wegener *et al.*, 2003).
- Ground meat is frequently associated with outbreaks of salmonellosis (Stock and Stolle, 2001).
- Up to 70% of foodborne illnesses are estimated to be linked to catered food (Filion and Powell, 2011; Hensen *et al.*, 2006; Jones *et al.*, 2004; Lee and Middleton, 2003).
- In Denmark, 61 of 86 reported outbreaks in 2011 were associated with outside-the-home settings (anonymous, 2012).
- To model the distribution of pathogens during the processing operation are of major relevance to risk analysts (Flores, 2006).


# Objectives of the study



Challenging  
Food safety

Quantitative Microbiological Risk Assessment - QMRA

risk of salmonellosis by  
consumption of "frikadeller",  
processed by the catering sector



Raw pork

Catering  
processing

Final meal




*Frikadelle*

Based on scenario analysis:

- to evaluate existing practices
- challenge the efficiency of different Food Safety Authorities recommendations

# Summarizing the performed work

Experimental work and modelling activities were performed:



Journal of  
Applied Microbiology



Journal of Applied Microbiology ISSN 1364-5072

ORIGINAL ARTICLE

## Modelling transfer of *Salmonella* Typhimurium DT104 during simulation of grinding of pork

C.O.A. Møller<sup>1</sup>, M.J. Nauta<sup>1</sup>, B.B. Christensen<sup>2</sup>, P. Dalgaard<sup>3</sup> and T.B. Hansen<sup>1</sup>

<sup>1</sup> National Food Institute, Technical University of Denmark, Søborg, Denmark

<sup>2</sup> Department of Food Science, Faculty of Life Sciences, University of Copenhagen, Frederiksberg C, Copenhagen, Denmark

<sup>3</sup> National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

Journal of Applied Microbiology

Volume 112, Issue 1, January 2012, Pages: 90–98



Contents lists available at SciVerse ScienceDirect

Food Microbiology

journal homepage: [www.elsevier.com/locate/fm](http://www.elsevier.com/locate/fm)



Effect of natural microbiota on growth of *Salmonella* spp. in fresh pork – A predictive microbiology approach

C.O.A. Møller<sup>a</sup>, Y. Ilg<sup>b</sup>, S. Aabo<sup>a</sup>, B.B. Christensen<sup>a,1</sup>, P. Dalgaard<sup>c</sup>, T.B. Hansen<sup>a,\*</sup>

<sup>a</sup> National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark

<sup>b</sup> Institute of Animal Science, University of Bonn, Katzenburgweg 7-9, 53115 Bonn, Germany


<sup>c</sup> National Food Institute, Technical University of Denmark, Søtofts Plads, Building 221, DK-2800 Kgs. Lyngby, Denmark

Food microbiology

Volume 34, Issue 2, June 2013, pages: 284-295



## Experimental work




- Process flow
- Time and temperature profiles
- Weight of ingredients
- Unit changes during the processing, e.g.:


1 kg meat pieces



200 g slices



200 g portions




5 kg batter



70-80 g  
*"frikadelle" batter*





## Risk of salmonellosis from consumption of “frikadeller”


### Modelling activities

Pathway

plus  
→

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study
Serving time plus cold storage until 6° C is reached	7. growth	Møller et al. (2013), this study
Estimation of the risk	9. dose response	FAO/WHO (2002)





## Risk of salmonellosis from consumption of 'frikadeller'

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study
Serving time plus cold storage until 6° C is reached	7. growth	Møller et al. (2013), this study
Estimation of the risk	9. dose response	FAO/WHO (2002)

plus  
→

2

QMRA  
framework

## Risk of salmonellosis from consumption of 'frikadeller'

Processing step	Basic activity	Source/model reference
-----------------	----------------	------------------------

Reception

1. initial contamination

**Hansen et al. (2010)**
 $R_{\text{prevalence}} = 4.2\%$  of the samples

39 positive samples within the interval 0.04–0.4 CFU/g

7 positive samples within the interval 0.4–4 CFU/g

5 positive samples within the interval 4–40 CFU/g

1 positive sample with more than 40 CFU/g (assumed max. 400 CFU/g)

~Histograma ( $\log(0.04)$ ,  $\log(400)$ , {39,7,5,1})

Lot 1

Pieces of pork - PC ( $1.289 \pm 109$  g)

PC1

PC2

PC3

PC4

PC5

1.  
Reception

Lot 2

Pieces of pork - PC ( $1.289 \pm 109$  g)

PC6

PC7

PC8

PC9

PC10


2

QMRA  
framework

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)

$$S_{\text{conc slice}} \sim \text{Multinomial}(\text{Rconc piece}, \{1/S_n \text{ slices}\})$$



2

QMRA  
framework

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)

$$M_i = (1-a_1)(1-a_2)(1-c_2) S_i + (b_1 \text{ gr}_{1,i-1}) + (b_2 \text{ gr}_{2,i-1})$$

Lot 1

Pieces of pork - PC ( $1.289 \pm 109$  g)

PC1

PC2

PC3

PC4

PC5

Slice 1

Slice 2

Slice 3

Slice 4

Slice 5

Portion 1

Portion 2

Portion 3

Portion 4

Portion 5

1.  
Reception2.  
Slicing  
(partitioning)3.  
Grinding slices  
into portions  
(cross-contamination)

Lot 2

Pieces of pork - PC ( $1.289 \pm 109$  g)

PC6

PC7

PC8

PC9

PC10

Slice 1

Slice 2

Slice 3

Slice 4

Slice 5

Portion 1

Portion 2

Portion 3


Portion 4

Portion 5

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)


- the meat typically corresponds to 2/3 of the total weight,
- so the total weight is the **weight of the meat multiplied by 3/2**.



## Risk of salmonellosis from consumption of “frikadeller”


Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)

~Multinomial( $M_{\text{conc batter lot}}, \{1/P_{\text{weight Meatball of } P_n \text{ MB samples}}\})$



## Risk of salmonellosis from consumption of “frikadeller”


Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study




## Heating inactivation in pan

Literature data	T (°C)	D (min)	log D
MURPHY et al., 2004.	55	45.87	1.66
	57.5	26.76	1.43
ground pork (40.2% fat)	60	5.07	0.71
	62.5	2.56	0.41
	65	1.91	0.28
	67.5	0.36	-0.44
	70	0.083	-1.08
JUNEJA et al., 2001	58	6.68	0.83
	60	6.65	0.83
Pork (8.5% fat)	62.5	1.62	0.21
	65	0.87	-0.06
JUNEJA et al., 2001	58	8.65	0.94
	60	5.48	0.74
Beef (12.5% fat)	62.5	1.5	0.18
	65	0.67	-0.17
SMITH et al., 2001.	55	9.05	0.96
	58	2.26	0.35
Low fat ground beef (4.8%)	61	0.57	-0.24
	64	0.15	-0.82
SMITH et al., 2001.	55	10.55	1.02
	58	2.15	0.33
Low fat ground beef (4.8%)	61	0.41	-0.39
	64	0.07	-1.15
SMITH et al., 2001.	55	10.27	1.01
	58	2.06	0.31
Low fat ground beef (4.8%)	61	0.43	-0.37
	64	0.14	-0.85
Velasquez et al. (2010)	55	3.846154	0.59
	58	0.653595	-0.18
Ground pork (2.5% fat)	60	0.263158	-0.58
	62	0.20284	-0.69
	63	0.119474	-0.92
JUNEJA et al., 2010	60	2.12766	0.33
Lean ground beef	65	0.364964	-0.44
	71	0.146628	-0.83

Salmonella sp in ground pork and S. Typhimurium DT104  
in low fat ground beef



five datapoints in the heating curve





# Heating inactivation in pan

**Appendix B** Observed data used to model heat inactivation of *Salmonella* during pan-frying.

Observed Batch	Start temperature <sup>a</sup> (°C)	Heating time (min)	End temperature (°C) per measured Meatball (MB)									
			MB 1	MB 2	MB 3	MB 4	MB 5	MB 6	MB 7	MB 8	MB 9	MB 10
2	5.1	7	19.9	23.5	31.9	34.2	39.1	39.5	39.8	42.7	44.5	56.3
3	5.1	7	18.2	18.3	20.1	20.6	27.0	28.3	29.5	41.0		
4	5.1	7	23.3	32.5	34.0	35.5	37.9	42.4	47.9	52.1	54.0	59.0
5	5.6	10	42.1	44.3	48.6	52.2	54.3	54.6				
6	7.2	10	37.2	46.0	49.1	50.1	62.2	66.8				
7	7.2	10	39.9	40.1	42.3	44.8	50	51.5				
8	8.4	6	32.1	33.1	39.1	53.2	55.2	59				
9	10.2	11	21.1	24.5	27.3	33.1	36.3	41.7	47			
10	10.2	5	45.1									

<sup>a</sup> Measured in the batter.



## Model approach

### Temperature profile:

$$T(t) = T_o + (T_{\text{end}} - T_o) / t$$

$$EF_{T_{\text{ref}}} = \sum_{\text{start}}^{\text{end}} 10^{\frac{T(t) - T_{\text{ref}}}{z}} \Delta t$$

$$LR_{\text{pan-frying}} = \frac{EF_{T_{\text{ref}}}}{D_{T_{\text{ref}}}}$$

## Parameter values

$$T_o = 5\text{-}10^{\circ} \text{ C}$$


$$T_{\text{end}} = 18\text{-}67^{\circ} \text{ C}$$

$$t = 7\text{-}11 \text{ min}$$

$$T_{\text{ref}} = 60^{\circ} \text{ C}$$

$$z = 7.34^{\circ} \text{ C}$$


$$D_{T_{\text{ref}}} = 1.8 \text{ min}$$



## Risk of salmonellosis from consumption of “frikadeller”


Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study

$$\sim \text{Poisson} (10^{(\log_{10}(\text{HP}_{\text{conc crit area}}) - 10^{(\text{HP}_{\log \log \text{reduction}}))})$$




## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study



## Heating inactivation in oven



Observed (●) and fitted with Newton's Law equation (—) temperature profiles of heating of meatball in oven until an end temperature of 95° C.

Model approach	Parameter values
Temperature profile:	$T_a = 121^{\circ}\text{C}$
$T(t) = T_a + (T_o - T_a) \cdot \exp(-k \cdot t)$	$T_o = 25\text{-}45^{\circ}\text{C}$
	$k = 6.1 \text{ h}^{-1}$
$EF_{Tref} = \Delta t$	$t = 4\text{-}11 \text{ min}$
$LR_{oven} = \frac{EF_{Tref}}{D_{Tref}}$	$T_{ref} = 65\text{-}75^{\circ}\text{C}$
	$z = 7.34^{\circ}\text{C}$
	$D_{Tref} = 0.4\text{-}0.02 \text{ min}$


### Data on eight meatballs used to determine the degree of heat inactivation in oven

Observed meatball	Start temperature ( $T_o$ , ° C) <sup>a</sup>	Heating time (t, min) <sup>b</sup>	End temperature (° C) <sup>c</sup>	$T_{ref}$ (° C)	Log ( $LR_{oven}$ ) <sup>d</sup>
1	25.4	11.1	89.8	95	3.74
		5.4	65.8	75	0.25
		5.2	64.6	74	0.07
		5.0	63.4	73	-0.10
		4.8	62.2	72	-0.27
		4.6	61.0	71	-0.44
		4.4	59.8	70	-0.61
		3.5	53.8	65	-1.46


## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study

~ Normal ( $HO_{avg \log \log red}$ ,  $HO_{st dev \log \log red}$ )




## Example of temperature profile for a meatball during processing at the catering sector




Involving: pan frying (grey full line), holding time (grey dashed line), heating in oven (grey dotted line), serving time (black full line), and chilled storage (black dotted line)

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study
Serving time plus cold storage until 6° C is reached	7. growth	Møller et al. (2013), this study



# Growth during storage



Observed (•, Δ) and fitted (grey line, black line) temperature profiles of the cooling of meatballs in kitchens one and two, respectively. Fitting done with Newton's Law resulting in  $k=1.8 \text{ h}^{-1}$  in both catering units.

## Model approach

Temperature profile:

$$T(t) = T_a + (T_o - T_a) \cdot \exp(-k \cdot t)$$

$$T_a = 20\text{-}30^\circ\text{C}$$

$$T_o = 18.2\text{-}66.8^\circ\text{C}$$

$$k = 1.8 \text{ h}^{-1}$$

$$t = 0\text{-}90 + 120 + 210 \text{ min}$$

Growth model:

$$\mu_{max} = (b \cdot (T - T_{min}) \cdot (1 - \exp(c \cdot (T - T_{max}))))^2$$

$$b = (0.04 \text{ h}^{-0.5} \cdot \text{C}^{-1}) \quad c = (0.43 \text{ C}^{-1})$$

$T$  is temperature


$$T_{min} (3.94^\circ \text{C})$$

$$T_{max} (48.0^\circ \text{C})$$

$$\text{Lag time} = RLT \cdot \ln(2) / \mu_{max}$$

where  $RLT$  is the relative lag time of 3.10

cooling (min)	Temperatur	umax (1/h)	lag (h)	lag (min)	coolingtid minus le	0=ikke vækst ; 1=vækst
12	0	38	1.61246	1.3	80	0
13	6	38	1.61246	1.3	80	0
14	12	38	1.61246	1.3	80	0
15	18	38	1.61246	1.3	80	0
16	24	38	1.61246	1.3	80	0
17	30	38	1.62437	1.3	79	0
18	36	37	1.50509	1.4	86	0
19	42	36	1.40895	1.5	91	0
20	48	35	1.33107	1.6	97	0
21	54	34	1.26772	1.7	102	0
22	60	33	1.21599	1.8	106	0
23	66	33	1.17361	1.8	110	0
24	72	32	1.13879	1.9	113	0
25	78	32	1.11010	1.9	116	0
26	84	32	1.08642	2.0	118	0
27	90	31	1.06684	2.0	121	0
28	96	31	1.05062	2.0	123	0
29	102	31	1.03716	2.1	124	0
30	108	31	1.02599	2.1	125	0
31	114	31	1.01671	2.1	127	0
32	120	31	1.00899	2.1	128	0
33	126	30	1.00256	2.1	128	0
34	132	30	0.99720	2.2	129	1
35	138	30	0.99274	2.2	130	1
36	144	30	0.98903	2.2	130	1
37	150	30	0.98593	2.2	131	1



54 min. growth of Salmonella at 30C

6 min. growth of Salmonella at 26C



## Growth during storage

**Table 4.** Change in log concentration  $\Delta \log C_x$  ( $T_o$ , RTK,  $t_{\text{holding}}$ ) for different start temperatures ( $T_o$ ) observed after pan-frying, for different room temperatures in the kitchen (RTK) and holding times  $t_{\text{holding}}$  between 0 and 90 min followed by 120 min serving time and cold storage at 6°C.

RTK – Room Temperature in the Kitchen (°C)	$T_o$	$\Delta \log C_x$ ( $T_o$ , RTK, $t_{\text{holding}}=90$ min)	Category <sup>a</sup>	$\Delta \log C_x$ ( $T_o$ , RTK, $t_{\text{holding}}=0$ min)	Category
30	18.2	0.72	2	0.00	2
	32.0	0.96	2	0.24	2
	33.0	1.00	2	0.28	2
	40.2	1.13	2	0.41	2
	42.0	1.17	2	0.46	2
	43.0	1.18	2	0.46	2
	44.0	1.22	2	0.50	2
	45.0	1.22	2	0.50	2
	46.0	1.23	2	0.51	2
	47.0	1.18	2	0.46	2
	55.0	0.58	3 <sup>b</sup>	-0.14	3
	56.0	0.17	3	-0.33	3
	57.0	-0.04	3	-0.64	3
	58.0	-0.26	3	-0.98	3
	59.0	-0.84	3	-1.57	3
	62.2	-3.64	3	-4.36	3
25	18.2	0.19	2	0.00	1
	32.0	0.40	2	0.00	1
	40.2	0.59	2	0.09	2
	46.0	0.64	2	0.17	2
	52.0	0.47	3 <sup>b</sup>	-0.01	3
	55.0	0.14	3	-0.33	3
	57.0	-0.26	3	-0.74	3
	59.9	-1.04	3	-1.50	3
	62.2	-3.40	3	-3.85	3
20	18.2	0.00	1	0.00	1
	32.0	0.00	1	0.00	1
	35.5	0.02	2	0.00	1
	40.2	0.15	2	0.00	1
	46.0	0.19	2	0.00	1
	53.0	-0.01	3	-0.19	3 <sup>c</sup>
	55.0	-0.22	3	-0.38	3
	57.0	-0.58	3	-0.74	3
	59.0	-1.27	3	-1.44	3
	62.2	-3.43	3	-3.58	3

<sup>a</sup> Process determining the result (see section 2.3.1.8). 1: lag phase; 2: growth; 3: inactivation.

<sup>b</sup> Both growth and inactivation can occur within a time temperature profile. The data point is assigned to the category where it gives the best fit to the linear model.

<sup>c</sup> No observations fall in category 2.


In the QMRA Monte Carlo  
Normal distribution of  $T_o$ :

- mean 40.3
- standard deviation 12

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
Holding time	7. growth	Møller et al. (2013), this study
Heat inactivation in oven	8. inactivation	this study
Serving time plus cold storage until 6° C is reached	7. growth	Møller et al. (2013), this study

$$= (10^{(\log_{10}(\text{HO}_{\text{conc after inac}}) + 10^{(\Delta \log C_x)))}$$



2

QMRA  
framework

## Risk of salmonellosis from consumption of “frikadeller”

Processing step	Basic activity	Source/model reference
Reception	1. initial contamination	Hansen et al. (2010)
Slicing	2. partitioning	Nauta (2005)
Grinding slices into portions	3. cross contamination	Møller et al. (2012)
Mixing of ingredients	4. mixing	Nauta (2005)
Dividing into meatballs	5. partitioning	Nauta (2005)
Heating inactivation in pan	6. inactivation	this study
— Holding time	7. growth	Møller et al. (2013), this study
plus Heat inactivation in oven	8. inactivation	this study
— Serving time plus cold storage until 6° C is reached	7. growth	Møller et al. (2013), this study
Estimation of the risk	9. dose response	FAO/WHO (2002)

$$Pill_{dose} \sim 1 - (1 + (C_{conc \text{ after } \Delta \log C_x} / \beta)^a$$


where  $C_{conc \text{ after } \Delta \log C_x}$  is the concentration of *Salmonella* spp. in a meatball at the time of serving,

$\alpha$  was -0.1324 and

$\beta$  equal to 51.45 (FAO/WHO, 2002).

The mean probability of illness from consumption of meatballs was used as the risk estimate

## Risk of salmonellosis from consumption of 'frikadeller'



# Baseline model

Processing step	Parameters
Reception	Prevalence = 4.2%
	Concentration = 91 % 0.04-4 CFU/g
	9 % → 400 CFU/g
Slicing	
Grinding slices into portions	
Mixing of ingredients	
Dividing into meatballs	
Heating inactivation in pan	
Holding time	
plus Heat inactivation in oven	75° C
→ Serving time plus cold storage until 6° C is reached	RTK = 20° C
Estimation of the risk	


6

## QMRA modelling


# Scenario analysis

- Evaluate processing practices
- Test control measures


### Scenario 1A – 1G



### Scenario 2 Baseline scenario RTK = 20° C



### Scenario 3A – 3C





# 6 QMRA modelling

## Challenge 1:

Are the recommendations of Food Safety Authorities good enough?

## Scenario analysis




6

QMRA  
modelling


# Scenario analysis

## Challenge 2:

Is it safe to  
consume cold  
“frikadeller”?



## Room Temperature in the Kitchen (RTK)




**Baseline scenario: no oven, RTK = 20° C**  
(1 case in 10,000 servings)

Tested scenarios

3C (no oven, rtk = 30°C)

3B (no oven, rtk = 25°C)

3A (no oven, rtk = 20°C)





6

QMRA  
modelling


# Scenario analysis

## Challenge 3:

How many people would become ill with salmonellosis in Denmark per year?

serving times/year = 24		Danish population (2013) = 5600000		20% of Danish population = 1120000				
1A (heating in oven = 75°C)	1B (heating in oven = 74°C)	1C (heating in oven = 73°C)	1D (heating in oven = 72°C)	1E (heating in oven = 71°C)	1F (heating in oven = 70°C)	1G (heating in oven = 65°C)	2 (no oven nor storage)	3C (no oven, rtk = 30°C)
0.000005	0.000012	0.000027	0.000044	0.000081	0.000122	0.000541	0.001217	0.003332
						5.41E-04	1.22E-03	3.33E-03
137	332	717	1186	2168	3291	14556	32706	89562

## Sensitivity analysis


Investigated aspect	Baseline model	Sensitivity analysis	Alternative scenario
✓ Fraction of "frikadelle" with Survival of <i>Salmonella</i>	 } = 33 %	<div><div>= 25 %</div><div>= 50 %</div></div>	<div><div>→ 4A</div><div>→ 4B</div></div>
✓ Heating inactivation according to meat matrix	→ z-value = 7.34	→ z-value = 5.90	→ 4C
✓ Prevalence of <i>Salmonella</i> in pork	→ 4.2 %	→ 20.0 %	→ 4D
✓ Concentration of <i>Salmonella</i> in pork	→ <div><div>91 % 0.04-4 CFU/g</div><div>9 % → 400 CFU/g</div></div>	→ <div><div>88 % 0.04-4 CFU/g</div><div>12 % 40-400 CFU/g</div></div>	→ 4E
✓ ↑ prevalence and ↑ concentration	→ 4F		

6

QMRA  
modelling

## Sensitivity analysis

scenario 1G, oven 65°C



↑ impact on the risk estimates



Combination of

↑ prevalence and ↑ concentration

# Remarks and future perspectives

## Risk assessment of *Salmonella* spp. in Danish meatballs produced at the catering sector

Cleide O. de A. Møller<sup>1</sup>, Maarten J. Nauta<sup>1</sup>, Donald W. Schaffner<sup>2</sup>, Paw Dalgaard<sup>3</sup>, Bjarke B. Christensen<sup>1,4</sup>, Tina B. Hansen<sup>1</sup>

- ✓ The model flexible structure allows scenario analysis
- ✓ Core temperatures  $> 70^{\circ}\text{C}$  inactivate *Salmonella*
- ✓ No growth of *Salmonella* if:
  - $RTK = < 20^{\circ}\text{C}$
  - For 3.5 h until refrigeration
  - Refrigerated storage =  $< 6^{\circ}\text{C}$


### To be investigated

- ✓ Heat inactivation :
  - Specific heating profile
  - “Frikadelle” batter
- ✓ Growth
  - Heat injured *Salmonella* cells
- ✓ Improvement of the previously developed models (transfer and growth)



# Acknowledgements

- ✓ This research project was financed by the Technical University of Denmark through the FoodDTU programme
- ✓ Constructive advice and critical comments were given by:



Tina Beck Hansen



Maarten Nauta



Donald W. Schaffner



Paw Dalgaard



Bjarke Bak Christensen



- ✓ Skillful technical assistance was provided by colleagues from the Division of Food Microbiology at the National Food Institute.



Rikke Krag



Mette Kemp



Kate Vibefeldt



Louise Vignæs